condition, such as a pimple or ingrown hair. Because of this, most people who have herpes do not know it.

Genital herpes sores usually appear as one or more blisters on or around the genitals, rectum or mouth. The blisters break and leave painful sores that may take weeks to heal. These symptoms are sometimes called "having an outbreak." The first time someone has an outbreak they may also have flu-like symptoms such as fever, body aches, or swollen glands.

Repeat outbreaks of genital herpes are common, especially during the first year after infection. Repeat outbreaks are usually shorter and less severe than the first outbreak. Although the infection can stay in the body for the rest of your life, the number of outbreaks tends to decrease over a period of years.

You should be examined by your doctor if you notice any of these symptoms or if your partner has an STD or symptoms of an STD, such as an unusual sore, a smelly discharge, burning when urinating, or, for women specifically, bleeding between periods.

How will my doctor know if I have herpes?

Often times, your healthcare provider can diagnose genital herpes by simply looking at your symptoms. Providers can also take a sample from the sore(s) and test it. Have an honest and open talk with your health care provider and ask whether you should be tested for herpes or other STDs.

Can herpes be cured?

There is no cure for herpes. However, there are medicines that can prevent or shorten outbreaks. One of these herpes medicines can be taken daily, and makes it less likely that you will pass the infection on to your sex partner(s).

What happens if I don't get treated?

Genital herpes can cause painful genital sores and can be severe in people with suppressed immune systems. If you touch your sores or the fluids from the sores, you may transfer herpes to another part of your body, such as your eyes. Do not touch the sores or fluids to avoid spreading herpes to another part of your body. If you touch the sores or fluids, immediately wash your hands thoroughly to help avoid spreading your infection.

Some people who get genital herpes have concerns about how it will impact their overall health, sex life, and relationships. It is best for you to talk to a health care provider about those concerns, but it also is important to recognize that while herpes is not curable, it can be managed. Since a genital herpes diagnosis may affect how you will feel about current or future sexual relationships, it is important to understand how to talk to sexual partners about STDs. You can find one resource here: GYT Campaign, http://www.cdcnpin.org/stdawareness/GYT.aspx

If you are pregnant, there can be problems for you and your unborn child. See "I'm pregnant. How could genital herpes affect my baby?" above for information about this.



Where can I get more information?

Division of STD Prevention (DSTDP)

Centers for Disease Control and Prevention

www.cdc.gov/std

Personal health inquiries and information about STDs:
CDC-INFO Contact Center
1-800-CDC-INFO (1-800-232-4636)
Contact www.cdc.gov/info

Resources:

CDC National Prevention Information Network (NPIN)

http://www.cdcnpin.org/scripts/index.asp

P.O. Box 6003 Rockville, MD 20849-6003 E-mail: info@cdcnpin.org

American Sexual Health Association (ASHA)

http://www.ashastd.org/

P.O. Box 13827 Research Triangle Park, NC 27709-3827 1-800-783-9877

Can I still have sex if I have herpes?

If you have herpes, you should tell your sex partner(s) and let him or her know that you do and the risk involved. Using condoms may help lower this risk but it will not get rid of the risk completely. Not having sores or other symptoms of herpes can also lower the risk, but again not completely. Even if you do not have any symptoms, you can still infect your sex partners.

What is the link between genital herpes and HIV?

Genital herpes can cause sores or breaks in the skin or lining of the mouth, vagina, and rectum. The genital sores caused by herpes can bleed easily. When the sores come into contact with the mouth, vagina, or rectum during sex, they increase the risk of giving or getting HIV if you or your partner has HIV.

Last reviewed: January 23, 2014 Page 2 of 2